

## 9.6

*What you should learn*

**GOAL 1** Solve a right triangle.

**GOAL 2** Use right triangles to solve **real-life** problems, such as finding the glide angle and altitude of a space shuttle in **Example 3**.

*Why you should learn it*

▼ To solve **real-life** problems such as determining the correct dimensions of a wheel-chair ramp in Exs. 39–41.



## Solving Right Triangles

**GOAL 1 SOLVING A RIGHT TRIANGLE**

Every right triangle has one right angle, two acute angles, one hypotenuse, and two legs. To **solve a right triangle** means to determine the measures of all six parts. You can solve a right triangle if you know either of the following:

- Two side lengths
- One side length and one acute angle measure

As you learned in Lesson 9.5, you can use the side lengths of a right triangle to find trigonometric ratios for the acute angles of the triangle. As you will see in this lesson, once you know the sine, the cosine, or the tangent of an acute angle, you can use a calculator to find the measure of the angle.

In general, for an acute angle  $A$ :

if  $\sin A = x$ , then  $\sin^{-1} x = m\angle A$ . ← The expression  $\sin^{-1} x$  is read as “the inverse sine of  $x$ .”

if  $\cos A = y$ , then  $\cos^{-1} y = m\angle A$ .

if  $\tan A = z$ , then  $\tan^{-1} z = m\angle A$ .

**ACTIVITY****Developing Concepts****Finding Angles in Right Triangles**

- 1** Carefully draw right  $\triangle ABC$  with side lengths of 3 centimeters, 4 centimeters, and 5 centimeters, as shown.


- 2** Use trigonometric ratios to find the sine, the cosine, and the tangent of  $\angle A$ . Express the ratios in decimal form.

- 3** In Step 2, you found that  $\sin A = \frac{3}{5} = 0.6$ . You can use a calculator to find  $\sin^{-1} 0.6$ . Most calculators use one of the keystroke sequences below.

$$\overbrace{\sin^{-1}}^{\sin^{-1}} \quad \boxed{2\text{nd}} \quad \boxed{\text{SIN}} \quad 0.6 \quad \boxed{\text{ENTER}} \quad \text{or} \quad 0.6 \quad \overbrace{\sin^{-1}}^{\sin^{-1}} \quad \boxed{2\text{nd}} \quad \boxed{\text{SIN}}$$


Make sure your calculator is in degree mode. Then use each of the trigonometric ratios you found in Step 2 to approximate the measure of  $\angle A$  to the nearest tenth of a degree.

- 4** Use a protractor to measure  $\angle A$ . How does the measured value compare with your calculated values?



**EXAMPLE 1** Solving a Right Triangle

Solve the right triangle. Round decimals to the nearest tenth.

**SOLUTION**

Begin by using the Pythagorean Theorem to find the length of the hypotenuse.

$$(\text{hypotenuse})^2 = (\text{leg})^2 + (\text{leg})^2$$

Pythagorean Theorem

$$c^2 = 3^2 + 2^2$$

Substitute.

$$c^2 = 13$$

Simplify.

$$c = \sqrt{13}$$

Find the positive square root.

$$c \approx 3.6$$

Use a calculator to approximate.

Then use a calculator to find the measure of  $\angle B$ :

$$\left( \left( 2 \div 3 \right) \right) \left[ 2^{\text{nd}} \right] \left[ \text{TAN} \right] \approx 33.7^\circ$$


Finally, because  $\angle A$  and  $\angle B$  are complements, you can write

$$m\angle A = 90^\circ - m\angle B \approx 90^\circ - 33.7^\circ = 56.3^\circ.$$

- The side lengths of the triangle are 2, 3, and  $\sqrt{13}$ , or about 3.6. The triangle has one right angle and two acute angles whose measures are about  $33.7^\circ$  and  $56.3^\circ$ .

**EXAMPLE 2** Solving a Right Triangle

Solve the right triangle. Round decimals to the nearest tenth.

**SOLUTION**

Use trigonometric ratios to find the values of  $g$  and  $h$ .

$$\sin H = \frac{\text{opp.}}{\text{hyp.}}$$

$$\cos H = \frac{\text{adj.}}{\text{hyp.}}$$

$$\sin 25^\circ = \frac{h}{13}$$

$$\cos 25^\circ = \frac{g}{13}$$

$$13 \sin 25^\circ = h$$

$$13 \cos 25^\circ = g$$

$$13(0.4226) \approx h$$

$$13(0.9063) \approx g$$

$$5.5 \approx h$$

$$11.8 \approx g$$

Because  $\angle H$  and  $\angle G$  are complements, you can write

$$m\angle G = 90^\circ - m\angle H = 90^\circ - 25^\circ = 65^\circ.$$

- The side lengths of the triangle are about 5.5, 11.8, and 13. The triangle has one right angle and two acute angles whose measures are  $65^\circ$  and  $25^\circ$ .

**STUDENT HELP****Study Tip**

There are other ways to find the side lengths in Examples 1 and 2. For instance, in Example 2, you can use a trigonometric ratio to find one side length, and then use the Pythagorean Theorem to find the other side length.

## GOAL 2 USING RIGHT TRIANGLES IN REAL LIFE

### EXAMPLE 3 Solving a Right Triangle

#### FOCUS ON CAREERS



#### ASTRONAUT

Some astronauts are pilots who are qualified to fly the space shuttle. Some shuttle astronauts are mission specialists whose responsibilities include conducting scientific experiments in space. All astronauts need to have a strong background in science and mathematics.




#### CAREER LINK

[www.mcdougallittell.com](http://www.mcdougallittell.com)

**SPACE SHUTTLE** During its approach to Earth, the space shuttle's glide angle changes.

- When the shuttle's altitude is about 15.7 miles, its horizontal distance to the runway is about 59 miles. What is its glide angle? Round your answer to the nearest tenth.
- When the space shuttle is 5 miles from the runway, its glide angle is about  $19^\circ$ . Find the shuttle's altitude at this point in its descent. Round your answer to the nearest tenth.



#### SOLUTION


- Sketch a right triangle to model the situation. Let  $x^\circ$  = the measure of the shuttle's glide angle. You can use the tangent ratio and a calculator to find the approximate value of  $x$ .

$$\tan x^\circ = \frac{\text{opp.}}{\text{adj.}}$$

$$\tan x^\circ = \frac{15.7}{59}$$

$$x = ( 15.7 \div 59 ) 2\text{nd TAN}$$

$$x \approx 14.9$$



Substitute.

Use a calculator to find  $\tan^{-1}\left(\frac{15.7}{59}\right)$ .

- When the space shuttle's altitude is about 15.7 miles, the glide angle is about  $14.9^\circ$ .

- Sketch a right triangle to model the situation. Let  $h$  = the altitude of the shuttle. You can use the tangent ratio and a calculator to find the approximate value of  $h$ .

$$\tan 19^\circ = \frac{\text{opp.}}{\text{adj.}}$$

$$\tan 19^\circ = \frac{h}{5}$$


$$0.3443 \approx \frac{h}{5}$$

$$1.7 \approx h$$

Substitute.

Use a calculator.

Multiply each side by 5.



- The shuttle's altitude is about 1.7 miles.

#### STUDENT HELP



#### HOMEWORK HELP

Visit our Web site [www.mcdougallittell.com](http://www.mcdougallittell.com) for extra examples.

## GUIDED PRACTICE

**Vocabulary Check** ✓

1. Explain what is meant by *solving* a right triangle.

**Concept Check** ✓

Tell whether the statement is **true** or **false**.

2. You can solve a right triangle if you are given the lengths of any two sides.  
3. You can solve a right triangle if you know only the measure of one acute angle.

**Skill Check** ✓




**CALCULATOR** In Exercises 4–7,  $\angle A$  is an acute angle. Use a calculator to approximate the measure of  $\angle A$  to the nearest tenth of a degree.


4.  $\tan A = 0.7$       5.  $\tan A = 5.4$       6.  $\sin A = 0.9$       7.  $\cos A = 0.1$

Solve the right triangle. Round decimals to the nearest tenth.


8.



9.



10.



## PRACTICE AND APPLICATIONS

### STUDENT HELP


**Extra Practice**  
to help you master  
skills is on p. 820.

**FINDING MEASUREMENTS** Use the diagram to find the indicated measurement. Round your answer to the nearest tenth.

11.  $QS$

12.  $m\angle Q$

13.  $m\angle S$




**CALCULATOR** In Exercises 14–21,  $\angle A$  is an acute angle. Use a calculator to approximate the measure of  $\angle A$  to the nearest tenth of a degree.


14.  $\tan A = 0.5$       15.  $\tan A = 1.0$       16.  $\sin A = 0.5$       17.  $\sin A = 0.35$   
18.  $\cos A = 0.15$       19.  $\cos A = 0.64$       20.  $\tan A = 2.2$       21.  $\sin A = 0.11$

**SOLVING RIGHT TRIANGLES** Solve the right triangle. Round decimals to the nearest tenth.

22.




23.




24.




25.



26.



27.



### STUDENT HELP

#### HOMEWORK HELP

**Example 1:** Exs. 11–27,  
34–37  
**Example 2:** Exs. 28–33  
**Example 3:** Exs. 38–41